

About the Celestial Objects

Listed on this page are several of the brighter, more interesting celestial objects visible in the evening sky this month (refer to the monthly sky map). The objects are grouped into three categories. Those that can be easily seen with the naked eye (that is, without optical aid), those easily seen with binoculars, and those requiring a telescope to be appreciated. **Note, all of the objects (except single stars) will appear more impressive when viewed through a telescope or very large binoculars.** They are grouped in this way to highlight objects that can be seen using the optical equipment that may be available to the star gazer.

Tips for Observing the Night Sky

When observing the night sky, and in particular deep-sky objects such as star clusters, nebulae, and galaxies, it's always best to observe from a dark location. Avoid direct light from street lights and other sources. If possible observe from a dark location away from the light pollution that surrounds many of today's large cities.

You will see more stars after your eyes adapt to the darkness—usually about 10 to 20 minutes after you go outside. Also, if you need to use a torch to view the sky map, cover the light bulb with red cellophane. This will preserve your dark vision.

Finally, even though the Moon is one of the most stunning objects to view through a telescope, its light is so bright that it brightens the sky and makes many of the fainter objects very difficult to see. So try to observe the evening sky on moonless nights around either New Moon or Last Quarter.

Astronomical Glossary

Conjunction – An alignment of two celestial bodies such that they present the least angular separation as viewed from Earth.

Constellation – A defined area of the sky containing a star pattern.

Diffuse Nebula – A cloud of gas illuminated by nearby stars.

Double Star – Two stars that appear close to each other in the sky; either linked by gravity so that they orbit each other (binary star) or lying at different distances from Earth (optical double). Apparent separation of stars is given in seconds of arc (").

Ecliptic – The path of the Sun's center on the celestial sphere as seen from Earth.

Elongation – The angular separation of two celestial bodies. For Mercury and Venus the greatest elongation occurs when they are at their most angular distance from the Sun as viewed from Earth.

Galaxy – A mass of up to several billion stars held together by gravity.

Globular Star Cluster – A ball-shaped group of several thousand old stars.

Light Year (ly) – The distance a beam of light travels at 300,000 km/sec in one year.

Magnitude – The brightness of a celestial object as it appears in the sky.

Open Star Cluster – A group of tens or hundreds of relatively young stars.

Opposition – When a celestial body is opposite the Sun in the sky.

Planetary Nebula - The remnants of a shell of gas blown off by a star.

Universal Time (UT) – A time system used by astronomers. Also known as Greenwich Mean Time. Singapore Standard Time is UT plus 8 hours.

Variable Star - A star that changes brightness over a period of time.

EQUATORIAL EDITION MAY 2024

Easily Seen with the Naked Eye

Arcturus	Boo	 Orange, giant K star. Name means "bear watcher". Dist=36.7 ly.
Procyon	CMi	• Greek name meaning "before the dog" rises before Sirius (northern latitudes). Dist=11.4 ly.
β Centauri	Cen	 With Alpha Centauri, forms the so-called "Pointers-to-the-Cross". Dist=525 ly.
α Centauri	Cen	 Nearest bright star to Sun at 4.4 ly. Brilliant double star in a telescope. 80 year period.
Coalsack	Cru	 Most famous naked-eye dark nebula. Requires dark sky. Dist=600 ly.
Castor	Gem	 Multiple star system with 6 components. 3 stars visible in telescope. Dist=52 ly.
Pollux	Gem	 With Castor, the twin sons of Leda in classical mythology. Dist=34 ly.
lpha Herculis	Her	Semi-regular variable. Magnitude varies between 3.1 & 3.9 over 90 days. Mag 5.4 companion.
Regulus	Leo	 Brightest star in Leo. A blue-white star with at least 1 companion. Dist=77 ly.
Antares	Sco	 Red, supergiant star. Name means "rival of Mars". Dist=135.9 ly.
Spica	Vir	• Latin name means "ear of wheat" and shown held in Virgo's left hand. Dist=250 ly.

Easily Seen with Binoculars

M44	Cnc	0	Praesepe or Beehive Cluster. Visible to the naked eye. Dist=590±20 ly.
M3	CVn	\oplus	Easy to find in binoculars. Might be glimpsed with the naked eye.
2808	Car	\oplus	Located 4 deg W of Nu Carinae. Visible to the naked eye on clear nights.
R Carinae	Car	•	Long period variable. Magnitude varies between 3.9 & 10.5 over 309 days.
3114	Car	0	Stunning open cluster. 30+ stars visible through 7x binoculars. Dist=2,900 ly.
3293	Car	0	Rich, tightly packed. Surrounded by large, faint nebulosity. Dist=8,500 ly.
IC 2602	Car	0	The "Five of Diamonds". Bright cluster twice diameter of full Moon. Dist=491 ly.
3372	Car		Eta Carinae Nebula. Enormous glowing cloud in rich star field. Dist=8,000 ly.
3532	Car	0	Herschel "most brilliant cluster". 60+ stars in 7x binoculars. Dist=1,300 ly.
ω Centauri	Cen	\oplus	Largest and brightest globular star cluster in sky. 1 million stars. Dist=17,000 ly.
Mel 111	Com	0	Coma Berenices. 80 mag 5-6 stars in 5 deg. Dist=288 ly. Age=400 million years.
4755	Cru	0	Jewel Box. Outstanding star cluster. Many contrasting colours. Dist=7,600 ly.
M13	Her	\oplus	Best globular in northern skies. Discovered by Halley in 1714. Dist=23,000 ly.
M48	Hya	0	12+ stars in 7x binoculars. Triangular asterism near centre. Dist=1,990 ly.
R Hydrae	Hya	•	Long period variable. Mag varies between 3.0 & 11.0 over 390 days. Brilliant red.
M12	0ph	\oplus	Close to the brighter M10. Dist=18,000 ly.
M10	0ph	\oplus	3 degrees from the fainter M12. Both may be glimpsed in binoculars. Dist=14,000 ly.
M4	Sco	\oplus	A close globular. May just be visible without optical aid. Dist=7,000 ly.
6231	Sco	0	Easy to see in binoculars. Dist=5,900 ly.
M5	Ser	\oplus	Fine globular star cluster. Telescope will reveal individual stars. Dist=25,000 ly.
6025	TrA	0	A small open star cluster in Milky Way. Dist=2,700 ly.
Mizar & Alcor	UMa	•	Good eyesight or binoculars reveals 2 stars. Not a binary. Mizar has a mag 4 companion.
IC 2391	Vel	()	Omicron Velorum Cluster. Superb object for binoculars. Dist=450 ly.

Telescopic Objects

γ Virginis

	ε Boötis	Boo	•	Red giant star (mag 2.5) with a blue-green mag 4.9 companion. Sep=2.8". Difficult to split.
	M67	Cnc	0	Contains 500+ stars mag 10 & fainter. One of the oldest clusters. Dist=2,350 ly.
	M94	CVn	0	Compact nearly face-on spiral galaxy. Dist=15 million ly.
	M51	CVn	0	Whirlpool Galaxy. First recognised to have spiral structure. Dist=25 million ly.
	3918	Cen	-	The Blue Planetary. Visible in a small telescope as a round blue disk.
	5128	Cen	0	Bisected by a wide obscuring lane. Strong radio source. Dist=11 million ly.
	M64	Com	0	Black-Eye Galaxy. Discovered by J.E. Bode in 1775 "a small, nebulous star".
	3242	Hya		Ghost of Jupiter. Bright blue disk. Mag 11 central star. Dist=2,600 ly.
	M83	Hya	0	Classic face-on spiral. Discovered in 1752 by Lacaille. In attractive star field.
	γ Leonis	Leo	•	Superb pair of golden-yellow giant stars. Mags 2.2 & 3.5. Orbit=600 years. Sep=4.4".
	5822	Lup	0	Large, attractive cluster. Dist=1,800 ly. Open cluster NGC 5823 to the south.
	k Puppis	Pup	•	Telescope easily shows two blue-white stars of almost equal brightness. Sep=9.9".
	6124	Sco	0	Contains 5 bright tightly packed stars near centre. 7 star chain. Dist=1,600 ly.
	M81	UMa	0	Beautiful spiral galaxy visible with binoculars. Easy to see in a telescope.
7	M82	UMa	0	Close to M81 but much fainter and smaller.
	3132	Vel		One of the brightest planetaries. Magnitude 10 central star. Dist=2,600 ly.
	M87	Vir	0	Supergiant galaxy with supermassive black hole at its core. Dist=53.5 million ly.
	M104	Vir	0	Sombrero Galaxy. Almost edge-on spiral galaxy. Protruding central core.

• Superb pair of mag 3.5 yellow-white stars. Orbit=169 years. At their closest in 2005.