

About the Celestial Objects

Listed on this page are several of the brighter, more interesting celestial objects visible in the evening sky this month (refer to the monthly sky map). The objects are grouped into three categories. Those that can be easily seen with the naked eye (that is, without optical aid), those easily seen with binoculars, and those requiring a telescope to be appreciated. Note, all of the objects (except single stars) will appear more impressive when viewed through a telescope or very large binoculars. They are grouped in this way to highlight objects that can be seen using the optical equipment that may be available to the star gazer.

Tips for Observing the Night Sky

When observing the night sky, and in particular deep-sky objects such as star clusters, nebulae, and galaxies, it's always best to observe from a dark location. Avoid direct light from street lights and other sources. If possible observe from a dark location away from the light pollution that surrounds many of today's large cities.

You will see more stars after your eyes adapt to the darkness—usually about 10 to 20 minutes after you go outside. Also, if you need to use a torch to view the sky map, cover the light bulb with red cellophane. This will preserve your dark vision.

Finally, even though the Moon is one of the most stunning objects to view through a telescope, its light is so bright that it brightens the sky and makes many of the fainter objects very difficult to see. So try to observe the evening sky on moonless nights around either New Moon or Last Quarter.

Astronomical Glossary

Conjunction – An alignment of two celestial bodies such that they present the least angular separation as viewed from Earth.

Constellation – A defined area of the sky containing a star pattern.

Diffuse Nebula – A cloud of gas illuminated by nearby stars.

Double Star – Two stars that appear close to each other in the sky; either linked by gravity so that they orbit each other (binary star) or lying at different distances from Earth (optical double). Apparent separation of stars is given in seconds of arc (").

Ecliptic – The path of the Sun's center on the celestial sphere as seen from Earth.

Elongation – The angular separation of two celestial bodies. For Mercury and Venus the greatest elongation occurs when they are at their most angular distance from the Sun as viewed from Earth.

Galaxy – A mass of up to several billion stars held together by gravity.

Globular Star Cluster – A ball-shaped group of several thousand old stars.

Light Year (ly) – The distance a beam of light travels at 300,000 km/sec in one year.

Magnitude – The brightness of a celestial object as it appears in the sky.

Open Star Cluster – A group of tens or hundreds of relatively young stars.

Opposition – When a celestial body is opposite the Sun in the sky.

Planetary Nebula – The remnants of a shell of gas blown off by a star.

Universal Time (UT) – A time system used by astronomers. Also known as Greenwich Mean Time. Singapore Standard Time is UT plus 8 hours.

Variable Star – A star that changes brightness over a period of time.

EQUATORIAL EDITION SEPTEMBER 2025

Easily Seen with the Naked Eye

Altaır	Aql	 Brightest star in Aquila. Name means "the flying eagle". Dist=16.8 ly.
lpha Centauri	Cen	• Nearest bright star to Sun at 4.4 ly. Brilliant double star in a telescope. 80 year period.
δ Cephei	Сер	Cepheid prototype. Mag varies between 3.5 & 4.4 over 5.366 days. Mag 6 companion.
Deneb	Cyg	 Brightest star in Cygnus. One of the greatest known supergiants. Dist=1,400±200 ly.
lpha Herculis	Her	Semi-regular variable. Magnitude varies between 3.1 & 3.9 over 90 days. Mag 5.4 companion.
Vega	Lyr	• The 5th brightest star in the sky. A blue-white star. Dist=25.0 ly.
Fomalhaut	PsA	 Brightest star in Piscis Austrinus. In Arabic the "fish's mouth". Dist=25 ly.
Antares	Sco	 Red, supergiant star. Name means "rival of Mars". Dist=135.9 ly.

Easily Seen with Binoculars

M31	And	0	The Andromeda Galaxy. Most distant object visible to naked eye. Dist=2.5 million ly.
M2	Aqr	\oplus	Resembles a fuzzy star in binoculars.
η Aquilae	Aql	•	Bright Cepheid variable. Mag varies between 3.6 & 4.5 over 7.166 days. Dist=1,200 ly.
6397	Ara	Φ	Thought to be the nearest globular. Dist=7,000 ly.
μ Cephei	Сер	•	Herschel's Garnet Star. One of the reddest stars. Mag 3.4 to 5.1 over 730 days.
χ Cygni	Cyg	•	Long period pulsating red giant. Magnitude varies between 3.3 & 14.2 over 407 days.
M39	Cyg	0	May be visible to the naked eye under good conditions. Dist=900 ly.
ν Draconis	Dra	•	Wide pair of white stars. One of the finest binocular pairs in the sky. Dist=100 ly.
M13	Her	\oplus	Best globular in northern skies. Discovered by Halley in 1714. Dist=23,000 ly.
M92	Her	\oplus	Fainter and smaller than M13. Use a telescope to resolve its stars.
ε Lyrae	Lyr	•	Famous Double Double. Binoculars show a double star. High power reveals each a double.
R Lyrae	Lyr	•	Semi-regular variable. Magnitude varies between 3.9 & 5.0 over 46.0 days.
M12	0ph	\oplus	Close to the brighter M10. Dist=18,000 ly.
M10	0ph	\oplus	3 degrees from the fainter M12. Both may be glimpsed in binoculars. Dist=14,000 ly.
IC 4665	0ph	0	Large, scattered open cluster. Visible with binoculars.
6633	0ph	0	Scattered open cluster. Visible with binoculars.
κ Pavonis	Pav	•	Cepheid-type. Magnitude varies between 3.9 & 4.8 over 9.088 days.
6752	Pav	\oplus	One of the better globular star clusters in the sky. Dist=14,000 ly.
M15	Peg	\oplus	Only globular known to contain a planetary nebula (Mag 14, d=1"). Dist=30,000 ly.
M8	Sgr		Lagoon Nebula. Bright nebula bisected by a dark lane. Dist=5,200 ly.
M25	Sgr	0	Bright cluster located about 6 deg N of "teapot's" lid. Dist=1,900 ly.
M22	Sgr	\oplus	A spectacular globular star cluster. Telescope will show stars. Dist=10,000 ly.
M4	Sco	\oplus	A close globular. May just be visible without optical aid. Dist=7,000 ly.
6231	Sco	0	Easy to see in binoculars. Dist=5,900 ly.
M6	Sco	0	Butterfly Cluster. 30+ stars in 7x binoculars. Dist=1,960 ly.
M7	Sco	Q.	Superb open cluster. Visible to the naked eye. Age=260 million years. Dist=780 ly.
M5	Ser	\oplus	Fine globular star cluster. Telescope will reveal individual stars. Dist=25,000 ly.
6025	TrA	0	A small open star cluster in Milky Way. Dist=2,700 ly.

Telescopic Objects

SMC

M11

M16

M27

Cr 399

reiesco	pic (JD	jects
7009	Aqr	÷	Saturn Nebula. Requires 8-inch telescope to see Saturn-like appendages.
7293	Aqr		Helix Nebula. Spans nearly 1/4 deg. Requires dark sky. Dist=300 ly.
Albireo	Cyg	•	Beautiful double star. Contrasting colours of orange and blue-green. Sep=34.4".
61 Cygni	Cyg	•	Attractive double star. Mags 5.2 & 6.1 orange dwarfs. Dist=11.4 ly. Sep=28.4".
γ Delphini	Del	•	Appear yellow & white. Mags 4.3 & 5.2. Dist=100 ly. Struve 2725 double in same field.
β Lyrae	Lyr	⊚	Eclipsing binary. Mag varies between 3.3 & 4.3 over 12.940 days. Fainter mag 7.2 blue star.
M57	Lyr		Ring Nebula. Magnificent object. Smoke-ring shape. Dist=4,100 ly.
M23	Sgr	0	Elongated star cluster. Telescope required to show stars. Dist=2,100 ly.
M20	Sgr		Trifid Nebula. A telescope shows 3 dust lanes trisecting nebula. Dist=5,200 ly.
M21	Sgr	0	A fine and impressive cluster. Dist=4,200 ly.
M17	Sgr		Omega Nebula. Contains the star cluster NGC 6618. Dist=4,900 ly.
6124	Sco	O	Contains 5 bright tightly packed stars near centre. 7 star chain. Dist=1,600 ly.

□ Eagle Nebula. Requires a telescope of large aperture. Dist=8,150 ly.

The Evening Sky Map (ISSN 1839-7735) Copyright © 2000–2025 Kym Thalassoudis. All Rights Reserved.

Wild Duck Cluster. Resembles a globular through binoculars. V-shaped. Dist=5,600 ly.

Dumbbell Nebula. Large, twin-lobed shape. Most spectacular planetary. Dist=975 ly.

Small Magellanic Cloud. Companion galaxy to Milky Way. Requires dark sky. Dist=210,000 ly.

Coathanger asterism or "Brocchi's Cluster". Not a true star cluster. Dist=218 to 1,140 ly.